

CONJUGACY, ORBIT EQUIVALENCE AND CLASSIFICATION OF MEASURE PRESERVING GROUP ACTIONS

Asger Törnquist

March 18, 2008

Abstract

We prove that if G is a countable discrete group with property (T) over an infinite subgroup $H \leq G$ which contains an infinite Abelian subgroup or is normal, then G has continuum many orbit inequivalent measure preserving a.e. free ergodic actions on a standard Borel probability space. Further, we obtain that the measure preserving a.e. free ergodic actions of such a G cannot be classified up to orbit equivalence by a reasonable assignment of countable structures as complete invariants. We also obtain a strengthening and a new proof of a non-classification result of Foreman and Weiss for conjugacy of measure preserving ergodic, a.e. free actions of discrete countable groups.

§1. INTRODUCTION

(A) Let G_1 and G_2 be discrete countable groups, acting by measure preserving transformations on standard Borel probability spaces (X, μ) and (Y, ν) , respectively, and giving rise to orbit equivalence relations E_{G_1} and E_{G_2} . Recall that the actions of G_1 and G_2 are *orbit equivalent* if there is a measure preserving transformation $T : X \rightarrow Y$ such that

$$xE_{G_1}y \iff T(x)E_{G_2}T(y).$$

The notion of orbit equivalence was introduced by Dye in [6],[7], who proved that any two ergodic $\mathbb{Z}$ -actions are orbit equivalent. Connes, Feldman and Weiss generalized this in [4] to all countable amenable groups, and eventually the combined work of Connes, Weiss [5], Schmidt [21], and Hjorth

[13] showed that this characterizes amenability: Any non-amenable discrete group has at least two non-orbit equivalent ergodic a.e. free measure preserving actions on a standard Borel probability space.

Recall that a group G is said to have the *relative property (T)* over a subgroup $H \leq G$ if there is $Q \subseteq G$ finite and $\varepsilon > 0$ such that whenever $\pi : G \rightarrow \mathcal{H}$ is a unitary representation of G on a Hilbert space $\mathcal{H}$ with a unit vector $\xi \in \mathcal{H}$ such that for all $g \in Q$

$$\|\pi(g)(\xi) - \xi\|_{\mathcal{H}} < \varepsilon,$$

then there is a non-zero H -invariant vector in π . The purpose of this paper is to prove the following (Theorem 4, §4):

1.1. THEOREM. *If G is a countable discrete group with the relative property (T) over an infinite subgroup $H \leq G$ such that either*

- (i) H contains an infinite Abelian subgroup or*
- (ii) H is normal in G ,*

then G has continuum many orbit inequivalent a.e. free measure preserving actions on a standard Borel probability space.

Such groups include well known examples such as the pair $(SL_2(\mathbb{Z}) \ltimes \mathbb{Z}^2, \mathbb{Z}^2)$, as well as any group with Property (T) outright, and groups that are formed by operations that preserve the classes (i) and (ii), such as free and direct products of arbitrary countable groups with a group with the relative property (T) over an infinite abelian subgroup.

It should be noted that (ii) in Theorem 1.1 was proven by Popa in [19]. We obtain a different proof here. The condition of normality can be weakened, see §3 Theorem 3.

(B) In descriptive set theory, a theory of classification has emerged over the last two decades, centered around the notion of *Borel reducibility*. If X and Y are Polish spaces and E and F are equivalence relations on X and Y , respectively, we say that E is Borel reducible to F if there is a Borel $f : X \rightarrow Y$ such that

$$(\forall x, y \in X) xEy \iff f(x)Ff(y).$$

We write $E \leq_B F$ if this is the case, and $E <_B F$ if $E \leq_B F$ but not $F \leq_B E$. Considering E and F as (potentially interesting) classification problems among the points of X and Y , respectively, we can interpret the

statement $E \leq_B F$ as saying that the points of X are classified up to E equivalence by a Borel assignment of F -equivalence classes as complete invariants. Alternatively, we may think of $E \leq_B F$ as an expression of the relative difficulty posed by the two classification problems. The reason that f is required to be Borel is to guarantee that the classification is reasonably “computable”, thus avoiding the useless classifications offered by the axiom of choice.

The notion of Borel reducibility will allow us to state the theorems of this paper in a much sharper way. We say that an equivalence relation E (on a Polish space X) is *smooth* or *concretely classifiable* if there is a Borel $f : X \rightarrow \mathbb{R}$ such that

$$(\forall x, y \in X) xEy \iff f(x) = f(y),$$

that is, if there is a complete classification using real numbers as invariants. A pivotal example of a *non-smooth* equivalence relation is E_0 , defined on $2^{\mathbb{N}}$ by

$$xE_0y \iff (\exists N)(\forall n \geq N) x(n) = y(n).$$

However, there are natural equivalence relations far beyond the complexity of E_0 . An important divide in the landscape of classification is the distinction between those equivalence relations that allow classification by *countable structures*, that is, allow a Borel assignment of invariants such as countable groups, graphs, trees, etc., considered up to the natural notion of isomorphism, as oppose to those that do not. (Precise definitions are given in §2 below.) Then we have the following sharper version of Theorem 1.1:

1.2. THEOREM. *If G is a countable discrete group with the relative property (T) over an infinite subgroup $H \leq G$ such that either*

- (i) H contains an infinite Abelian subgroup or*
- (ii) H is normal in G ,*

Then orbit equivalence of measure preserving ergodic G actions is not smooth and cannot be classified by countable structures.

It should be noted that by a result in [22], which was based on the result of Popa in [19], that in the case (ii) above it was known that the isomorphism relation for countable torsion free abelian groups $\simeq_{\text{TFA}_{\aleph_0}}$ is Borel reducible to orbit equivalence. Thus the above Theorem shows that it is in fact strictly above $\simeq_{\text{TFA}_{\aleph_0}}$ in the $\leq_B$ -hierarchy, and rules out that countable discrete groups can, in a reasonable way, be used as complete invariant for orbit equivalence of groups with the relative property (T).

We refer the reader to the introduction of [14] for further background on the very interesting subject of Borel reducibility and the descriptive set theory of classification problems.

(C) The proof of Theorem 1.2 (and 1.1) is based on the method used by Hjorth in [13], relating *conjugacy* of measure preserving actions to orbit equivalence. The most important case is when G is a group with the relative property (T) over an infinite Abelian subgroup $H \leq G$.

Let σ and τ be measure preserving G actions on a standard Borel probability space (X, μ) . We will say that σ and τ are *conjugate on H* if $\sigma|_H$ and $\tau|_H$ are conjugate. We write $\sigma \simeq_H \tau$. We say that σ is *ergodic on H* if $\sigma|_H$ is ergodic. In §4 we prove the following relativization of Corollary 2.6 in [13]:

1.3. LEMMA. *Let G be a countable group with the relative property (T) over the infinite subgroup $H \leq G$. Then at most countably many ergodic-on- H probability measure preserving actions of G that are not conjugate on H can be orbit equivalent.*

Given this, the task becomes to produce many G actions that are non-conjugate and ergodic on H . To this end we prove a version of Foreman and Weiss anti-classification result in [8] (Theorem A) for extensions of actions. This is based on Theorem 5.1 of Kechris and Sofronidis in [17]:

THEOREM (Kechris-Sofronidis). *Let X be an uncountable Polish space and let $P_c(X)$ be the Polish space of continuous Borel probability measures. Then absolute equivalence $\approx$ in $P_c(X)$ is generically turbulent¹ and thus it is not smooth and not classifiable by countable structures.*

Assuming now that $H \leq G$ is an infinite Abelian group, then for each symmetric measure σ on the dual $\hat{H}$ we have an associated real positive definite function

$$\varphi_\sigma(h) = \int \chi(h) d\sigma(\chi).$$

Then if $\bar{\varphi}_\sigma$ is the trivial extension to G , then we may construct a Gaussian action ρ_σ of G as in [9] p. 90-91. In Lemma 3.5 we will compute a formula for the maximal spectral type of the Koopman representation $\kappa^\sigma|_H$ of this action restricted to H . We then show that if the measure σ has support on a “sufficiently independent” set (see Definition 3.11), then $\sigma \mapsto \rho_\sigma$

¹See §2 for a definition.

provides us with a Borel function such that equivalent measures produce conjugate G -actions, and inequivalent measures produce G -actions that are non-conjugate on H . Furthermore, the action $\rho_\sigma|_H$ will be weakly mixing. Thus $\sigma \mapsto \rho_\sigma$ is a Borel reduction of $\approx$ to $\simeq_H$.

We note that we obtain a new proof and a stronger version of Foreman and Weiss' result (Theorem 1, 2 and 3, §3):

1.4. THEOREM. *Let G be a countable discrete group and let $H \leq G$ be an infinite subgroup. Let (X, μ) be a standard Borel probability space. Then:*

(i) The measure preserving H -actions on (X, μ) that can be extended to an ergodic measure preserving G -action on (X, μ) cannot be classified up to conjugacy by countable structures.

(ii) If H either contains an infinite Abelian subgroup or is normal in G , then the measure preserving ergodic H -actions on (X, μ) that can be extended to an ergodic measure preserving G -action on (X, μ) are not classifiable by countable structures.

(D) The author would like to thank Greg Hjorth for useful discussions about the ideas for this paper at an early stage while the author was visiting the Mathematics Department and Logic Center at UCLA, in July 2006, and Alexander Kechris for making available his extensive new notes on measure preserving actions [16], the appendices of which were indispensable to this work. The author would also like to thank the referee, whose comments and suggestions have improved this paper vastly.

Research for this paper was supported in part by the Danish Natural Science Research Council grant no. 272-06-0211.

§2. BOREL REDUCIBILITY

2.1. Notation: Throughout this paper, if E is an equivalence relation on some set X and $Y \subseteq X$, we will write E^Y for $E|_{Y \times Y}$.

2.2. Reducibility notions. We first introduce two variant technical notions of Borel reducibility that will be of useful for us in this paper. Let X, Y be Polish spaces, and let E and F be equivalence relations on X and

Y , respectively. Recall that we say that E is Borel reducible to F , written $E \leq_B F$, if there is a Borel $f : X \rightarrow Y$ such that

$$(\forall x, y \in X) xEy \iff f(x)Ff(y).$$

If F' is another equivalence relation on Y and $F \subseteq F'$ then it is natural to introduce the following stronger notion: We will say that E is Borel reducible to the pair (F, F') , written $E \leq_B (F, F')$, if there is a Borel function $f : X \rightarrow Y$ such that

$$(\forall x, y \in X) xEy \implies f(x)Ff(y) \wedge f(x)F'f(y) \implies xEy.$$

It is clear that if $E \leq_B (F, F')$ then for any equivalence relation $F \subseteq F'' \subseteq F'$ we have $E \leq_B F''$.

A useful weaker reducibility notion is the following: We say that E is *Borel countable-to-one reducible* to F , written $E \leq_B^\omega F$, if there is a Borel $f : X \rightarrow Y$ such that

$$xEy \implies f(x)Ff(y)$$

and the induced map $\hat{f} : X/E \rightarrow Y/F : \hat{f}([x]_E) = [f(x)]_F$ is countable-to-1.

2.3. Classification by countable structures. Recall that an equivalence relation E *admits a classification by countable structures* (or *is classifiable by countable structures*) if there is a countable language $\mathcal{L}$ such that $E \leq_{B \simeq_{\text{Mod}(\mathcal{L})}}$, where $\simeq_{\text{Mod}(\mathcal{L})}$ is isomorphism for countable models of $\mathcal{L}$ (see [14] for details.) Let E be an equivalence relation on a Polish space X . We will say that E is *generically turbulent* if

- i) All E classes are meagre and
- ii) There is a Polish group G acting continuously on X such that the induced orbit equivalence relation $E_G \subseteq E$ and the action of G on X is generically turbulent in the sense of [14].

Then it is immediate from [14], Theorem 3.18, and [1], Theorem 3.4.5, that if E is generically turbulent then $E_0 \leq_B E$ and E is not classifiable by countable structures. Further we have:

2.4. PROPOSITION. *Let F be an equivalence relation on a Polish space Y , and suppose there is a generically turbulent equivalence relation E on a Polish space X such that $E \leq_B^\omega F$. Then $E_0 <_B F$ and F does not admit a classification by countable structures.*

Proof. Let f witness that $E \leq_B^\omega F$. Define

$$xE'y \iff f(x)Ff(y).$$

Then $E \subseteq E'$ and E' has countable index over E . Thus E' is generically turbulent. Since by definition $E' \leq_B F$ it follows that $E_0 \leq_B F$ and that F is not classifiable by countable structures. $\square$

§3. CONJUGACY ON A SUBGROUP

Let (X, μ) be a standard Borel probability space. We denote by $\text{Aut}(X, \mu)$ the group of measure preserving transformations on X , which we give the weak topology (c.f. [15]). Further, if G is a countable group, we denote by $\mathcal{A}(G, X, \mu)$ the set of homomorphisms of G to $\text{Aut}(X, \mu)$ equipped with the subspace topology it inherits from $\text{Aut}(X, \mu)^G$, and identify $\mathcal{A}(G, X, \mu)$ with the space of measure preserving transformations. $\mathcal{E}(G, X, \mu)$ and $\mathcal{W}(G, X, \mu)$ denote the subsets of ergodic, respectively weakly mixing, measure preserving actions of G on X . We denote by $\mathcal{A}^*(G, X, \mu) \subseteq \mathcal{A}(G, X, \mu)$ subset of a.e. free actions, and we let

$$\begin{aligned}\mathcal{E}^*(G, X, \mu) &= \mathcal{E}(G, X, \mu) \cap \mathcal{A}^*(G, X, \mu) \\ \mathcal{W}^*(G, X, \mu) &= \mathcal{W}(G, X, \mu) \cap \mathcal{A}^*(G, X, \mu).\end{aligned}$$

$\text{Aut}(X, \mu)$ acts on $\mathcal{A}(G, X, \mu)$ by conjugation. We denote by $\simeq$ the conjugation relation in $\mathcal{A}(G, X, \mu)$.

If H is a subgroup of G , then we denote by $\mathcal{E}_H(G, X, \mu)$ the subset of $\sigma \in \mathcal{E}(G, X, \mu)$ such that $\sigma|_H$ is ergodic as an action of H , that is, the ergodic on H m.p. actions of G as defined in the introduction. Similarly, we denote by $\mathcal{W}_H(G, X, \mu)$ the set of $\sigma \in \mathcal{W}(G, X, \mu)$ such that $\sigma|_H$ is weakly mixing. We denote by $\simeq_H$ the equivalence relation in $\mathcal{A}(G, X, \mu)$ of being conjugate on H : That is, for $\sigma, \tau \in \mathcal{A}(G, X, \mu)$ we write $\sigma \simeq_H \tau$ iff there is $T \in \text{Aut}(X, \mu)$ such that on a measure one set we have

$$(\forall h \in H) T(\sigma(h)(x)) = \tau(h)(T(x)).$$

Finally, we define the corresponding sets of free actions $\mathcal{E}_H^*(G, X, \mu)$ and $\mathcal{W}_H^*(G, X, \mu)$ as before.

Let X be a Polish space. We will denote by $P(X)$ the Polish space of all Borel probability measures, c.f. [15] pp. 109. We denote by $P_c(X)$

the subset of continuous (i.e. non atomic) probability measures. If X is uncountable then $P_c(X)$ is a dense G_δ subset of $P(X)$, cf. [17]. Absolute equivalence $\approx$ is a Borel equivalence relation in $P(X)$, c.f. [15] 17.39. In this section we will prove the following:

THEOREM 1. *Let G be a countable group and $A \leq G$ an infinite abelian subgroup, and let (X, μ) be a non-atomic standard Borel probability space. Then $\approx^{P_c(2^\mathbb{N})} \leq_B (\simeq^{\mathcal{W}_A^*(G, X, \mu)}, \simeq_A^{\mathcal{W}_A^*(G, X, \mu)})$. So for any $A \leq H \leq G$ we have that $E_0 <_B \simeq_H^{\mathcal{W}_H^*(G, X, \mu)}$ and $\simeq_H^{\mathcal{W}_H^*(G, X, \mu)}$ is not classifiable by countable structures.*

The proof is presented in a sequence of lemmata. For the rest of this section, $A \leq G$ is a fixed infinite abelian subgroup of G , and G is a fixed countable group. We also fix once and for all a sequence (g_i) of coset representatives, and suppose that the identity is chosen to represent A .

The idea of the proof is the following: Given a finite (continuous) measure σ on the dual $\hat{A}$, we have the corresponding unitary π^σ on $L^2(\hat{A}, \sigma)$. If σ is a symmetric measure, then the corresponding positive definite function φ_σ is real valued. Considering then $\text{Ind}_A^G(\pi^\sigma)$ of G , which corresponds to the trivial extension $\bar{\varphi}_\sigma$ of φ_σ , we can then obtain a Gaussian G -action ρ_σ as in [9] (p. 90-91). Considering this action as an A -action, we will compute the maximal spectral type of the corresponding Koopman representation, using a notion of “lifted” spectral measure, introduced below. We will then show that if the measures σ, σ' have support on a suitably chosen perfect subset C of $\hat{A}$, then the resulting Gaussian action $\rho_\sigma|_A$ and $\rho_{\sigma'}|_A$ are conjugate as A -actions if and only if σ and σ' are absolutely equivalent.

3.1. Lifts. Let $H < A$ be a subgroup of the abelian group A . We define $P_H : \hat{A} \rightarrow \hat{H}$ by

$$P_H(\chi) = \chi|_H.$$

Let σ be a positive finite measure on $\hat{H}$, and let $\tau : \hat{H} \rightarrow P(\hat{A}) : \chi \mapsto \tau_\chi$ be a Borel assignment of probability measures such that for all $\chi \in \hat{H}$ we have

$$\tau_\chi(P_H^{-1}(\chi)) = 1.$$

Then we define the τ -lift of σ to be the measure σ_τ on $\hat{A}$ defined by

$$\int f d\sigma_\tau = \int \int f(\eta) d\tau_\chi(\eta) d\sigma(\chi).$$

Note that the lifting process respects absolute continuity: if $\sigma \ll \sigma'$ then $\sigma_\tau \ll \sigma'_\tau$. Note also that if $H' \leq H$ is a subgroup, then $P_{H'}[\sigma_\tau] = P_{H'}[\sigma]$. In particular, $P_H[\sigma_\tau] = \sigma$.

3.2. Haar lifts. A special instance of this construction arises as follows: Identify $\widehat{A/H}$ with the set of characters $\chi \in \hat{A}$ such that $\chi|_H = 1$, and let $\lambda_{\widehat{A/H}}$ be the (normalized) Haar measure on $\widehat{A/H}$. Then for $\chi \in \hat{H}$ define for a Borel $B \subseteq \hat{A}$,

$$\lambda_\chi(B) = \lambda_{\widehat{A/H}}(\tilde{\chi}^* B),$$

where $\tilde{\chi}$ is some (any) extension of χ to a character on $\hat{A}$.² Note that λ_χ is independent of the choice of extension $\tilde{\chi}$ since $\lambda_{\widehat{A/H}}$ is translation invariant. Then $\lambda_\chi(P_H^{-1}(\chi)) = 1$ and $\chi \mapsto \lambda_\chi$ is a Borel assignment of Borel probability measures on $\hat{A}$. For a measure σ on $\hat{H}$ we call the lifted measure σ_λ on $\hat{A}$ the *Haar lift* of σ , and denote it σ_H^A , with a purposely suggestive notation. It is routine to check that if $\varphi_\sigma : H \rightarrow \mathbb{C}$ is the positive definite function corresponding to the spectral measure σ on $\hat{H}$, then σ_H^A is the spectral measure of the trivial extension of φ_σ to A .

For each coset representative g_i , define $H_i = A \cap g_i A g_i^{-1}$ and $H'_i = g_i^{-1} A g_i \cap A$. We also define $\varphi_i : H_i \rightarrow H'_i$ to be the isomorphism

$$\varphi_i(a) = g_i^{-1} a g_i.$$

Note that φ_i only depends on the coset, not on the choice of representative. We denote by $\hat{\varphi}_i : \hat{H}'_i \rightarrow \hat{H}_i$ the associated isomorphism of the character groups, $\hat{\varphi}_i(\chi) = \chi \circ \varphi_i$. More generally, and in a slight abuse of notation, we will also denote by $\hat{\varphi}_i$ the homomorphism $\hat{A} \rightarrow \hat{H}_i$ defined by $\hat{\varphi}_i(\chi) = \chi \circ \varphi_i$. If there is any danger of confusion, that latter will be written $\hat{\varphi}_i \circ P_{H'_i}$. We leave the proof of the next Lemma as an easy exercise for the reader in using the definition of lifts and induced representation:

3.3. LEMMA. *Suppose $(\pi, \mathcal{X})$ is a unitary representation of A on a separable Hilbert space $\mathcal{X}$ and that A is a subgroup of some countable discrete group G . If the maximal spectral type of π is σ , then the maximal*

²We note that we may find a Borel $\chi \mapsto \tilde{\chi}$ function for this purpose since we may find a transversal for the equivalence relation induced by the natural action of the compact group $\widehat{A/H}$ on $\hat{A}$, using [16], Theorem 12.16.

spectral type of $\text{Ind}_A^G(\pi)|_A$ is

$$\sigma_{\text{Ind}_A^G(\pi)|_A}^{\max} \approx \sum_i \frac{1}{2^i} \hat{\varphi}_i[\sigma]_{H_i}^A.$$

3.4. Let σ be a symmetric, finite and positive measure on $\hat{A}$, so that the associated positive definite function φ_σ is real valued. Let $\bar{\varphi}_\sigma$ be the trivial extension of φ_σ to G , and let $\mu_\sigma = \mu_{\bar{\varphi}_\sigma}$ be the associated Gaussian measure on $\mathbb{R}^G$, which we denote ρ_σ . If κ is the associated Koopman representation on $L^2(\mathbb{R}^G, \mu)$ then

$$\kappa^{H_1^{(c)}(\mu)} \sim_u \text{Ind}_A^G(\pi^\sigma),$$

where $H_1^{(c)}$ is the 1st Wiener Chaos. It follows from Lemma 3.3 that the maximal spectral type of $\kappa^{H_1^{(c)}(\mu)}|_A$, which we denote by w_σ , is

$$w_\sigma = \sum_i \frac{1}{2^i} \hat{\varphi}_i[\sigma]_{H_i}^A.$$

Thus the maximal spectral type of the Koopman representation κ is $\exp(w_\sigma)$, from which we obtain:

3.5. LEMMA. *With notation as in 3.4, the maximal spectral type of $\kappa_0|_A$ is*

$$\sum_{\substack{i_1, \dots, i_n \in \mathbb{N} \\ n \in \mathbb{N}}} \frac{1}{2^n 2^{i_1 + \dots + i_n}} \hat{\varphi}_{i_1}[\sigma]_{H_{i_1}}^A * \dots * \hat{\varphi}_{i_n}[\sigma]_{H_{i_n}}^A.$$

3.6. Definition. For a finite positive measure σ on $\hat{A}$, define

$$\Phi_\sigma = \sum_{\substack{i_1, \dots, i_n \in \mathbb{N} \\ n \in \mathbb{N}}} \frac{1}{2^n 2^{i_1 + \dots + i_n}} \hat{\varphi}_{i_1}[\sigma]_{H_{i_1}}^A * \dots * \hat{\varphi}_{i_n}[\sigma]_{H_{i_n}}^A,$$

so that the previous Lemma states that the maximal spectral type of $\kappa_0|_A$ on $L_0^2(\mathbb{R}^G, \mu_\sigma)$ is Φ_σ .

The following Lemma is routine and we omit the proof.

3.7. LEMMA. Let $H < A$ be a subgroup of A with infinite index and suppose $C \subseteq \hat{A}$ is a Borel set such that for $P_H|C$ is 1-1. Then:

(i) For any finite positive Borel measure σ on $\hat{H}$ and $\chi \in \hat{A}$ it holds for the Haar lift that $\sigma_H^A(\chi C) = 0$.

(ii) For any finite positive Borel measure σ on $\hat{H}$ and any finite positive Borel measure τ on $\hat{A}$ we have that

$$\tau * \sigma_H^A(C) = 0.$$

3.8. Definition. A subset $C \subseteq \hat{A}$ is called *sufficiently independent* if

1. For all i such that H_i is infinite we have that $P_{H_i}|C$ is 1-1, and for all i such that H'_i is infinite we have that $P_{H'_i}|C$ is 1-1.
2. For all $\chi_1, \dots, \chi_k, \chi_{k+1} \in C$ distinct and all $H_{i_1}, \dots, H_{i_k}$ with $[A : H_{i_j}] < \infty$ for all $j = 1, \dots, k$, and all $s : \{1, \dots, k\} \rightarrow \{-1, 1\}$ we have that

$$\prod_{j=1}^k \chi_j^{s(j)} \circ \varphi_{i_j}|_{H_{(i_1, \dots, i_k)}} \neq \chi_{k+1}|_{H_{(i_1, \dots, i_k)}},$$

where

$$H_{(i_1, \dots, i_k)} = \bigcap_{j=1}^k H_{i_j}.$$

Moreover, for a measure σ on $\hat{A}$ we define the *symmetrization* $\hat{\sigma}$ of σ by

$$\hat{\sigma}(B) = \frac{1}{2}(\sigma(B) + \sigma(B^*)),$$

where $B \subseteq \hat{A}$ is Borel. Note that if $\psi : \hat{A} \rightarrow H$ is a Borel homomorphism of $\hat{A}$ into some (Polish) group H then $\psi[\hat{\sigma}] = \widehat{\psi[\sigma]}$.

The following is immediate from the definition of being sufficiently independent:

3.9. LEMMA. Suppose C is a sufficiently independent subset of $\hat{A}$. If σ is a continuous Borel measure on C then for any i such that H_i is infinite $P_{H_i}[\sigma]$ and $P_{H'_i}[\sigma]$ are continuous. Moreover, the measure $\hat{\varphi}_i[\sigma]$ is continuous.

We then have:

3.10. LEMMA. Suppose C is a perfect sufficiently independent subset of $\hat{A}$ such that $\lambda_{\hat{A}}(C) = 0$, and let σ, τ be finite non-atomic positive Borel measures on C . Then $\sigma \not\approx \tau$ implies $\Phi_{\hat{\sigma}} \not\approx \Phi_{\hat{\tau}}$.

Proof. Suppose $B \subseteq C$ is a Borel set such that $\sigma(B) > 0$ and $\tau(B) = 0$. Note that by definition $\sigma \ll \Phi_{\hat{\sigma}}$, thus $\Phi_{\hat{\sigma}}(B) > 0$. For simplicity, we will write σ_i for $\hat{\varphi}_i[\sigma]_{H_i}^A$ and τ_i for $\hat{\varphi}_i[\tau]_{H_i}^A$.

We claim that $\Phi_{\hat{\tau}}(B) = 0$. For this, it suffices to show that for all $i_1, \dots, i_n \in \mathbb{N}$ we have

$$\hat{\tau}_{i_1} * \dots * \hat{\tau}_{i_n}(B) = 0.$$

If for some $1 \leq k \leq n$ we have that $[A : H_{i_k}] = \infty$ and $|H_{i_k}| = \infty$ then it follows from Lemma 3.7 that

$$\hat{\tau}_{i_1} * \dots * \hat{\tau}_{i_n}(B) = 0.$$

Also, if $|H_{i_k}| < \infty$ for some $1 \leq k \leq n$ is clear that $\hat{\tau}_{i_k} \ll \lambda_{\hat{A}}$. It follows that³

$$\prod_{j=1}^n * \hat{\tau}_{i_j} \ll \left(\prod_{\substack{j=1 \\ j \neq k}}^n * \hat{\tau}_{i_j} \right) * \lambda_{\hat{A}} = \lambda_{\hat{A}},$$

where the latter follows from the translation invariance of the Haar measure $\lambda_{\hat{A}}$. Since we assumed $\lambda_{\hat{A}}(C) = 0$ it follows that

$$\hat{\tau}_{i_1} * \dots * \hat{\tau}_{i_n}(B) = 0.$$

So without loss of generality, assume that $[A : H_{i_j}] < \infty$ for all $j = 1, \dots, n$. Let

$$H = H_{(i_1, \dots, i_n)} = \bigcap_{j=1}^n H_{i_j}.$$

Then $[A : H] < \infty$, too. Thus P_H is finite-to-1 and so $P_H(B)$ is Borel. We will show that

$$P_H\left[\prod_{j=1}^n * \hat{\tau}_{i_j}\right](P_H(B)) = 0,$$

from which it clearly follows that $\hat{\tau}_{i_1} * \dots * \hat{\tau}_{i_n}(B) = 0$.

Since P_H is a homomorphism we have

$$P_H\left[\prod_{j=1}^n * \hat{\tau}_{i_j}\right] = \prod_{j=1}^n * P_H[\hat{\tau}_{i_j}].$$

³We use the notation $\prod_{i=1}^n * \mu_i = \mu_1 * \dots * \mu_n$.

Further, we have $P_H[\hat{\tau}_{i_j}] = P_H \circ \hat{\varphi}_{i_j}[\hat{\tau}]$ since $\hat{\tau}_{i_j} = \hat{\varphi}_{i_j}[\hat{\tau}]_{H_{i_j}}^A$.

Suppose for a contradiction that

$$P_H[\prod_{j=1}^n * \hat{\tau}_{i_j}](P_H(B)) = \prod_{j=1}^n * P_H \circ \hat{\varphi}_{i_j}[\hat{\tau}](P_H(B)) > 0.$$

By definition then

$$P_H \circ \hat{\varphi}_{i_1}[\hat{\tau}] \times \cdots \times P_H \circ \hat{\varphi}_{i_n}[\hat{\tau}](\{(\chi_1, \dots, \chi_n) \in \hat{H} : \chi_1 \cdots \chi_n \in P_H(B)\}) > 0.$$

It follows by repeated applications of Fubini's Theorem we may then find $\chi_1, \dots, \chi_n \in C \setminus B$ distinct and $s : \{1, \dots, n\} \rightarrow \{-1, 1\}$ such that

$$P_H \circ \hat{\varphi}_{i_1}(\chi_1^{s(1)}) \cdots P_H \circ \hat{\varphi}_{i_n}(\chi_n^{s(1)}) = P_H(\eta)$$

for some $\eta \in B$. But this says precisely that

$$\prod_{j=1}^n \chi_j^{s(j)} \circ \varphi_{i_j}|_H = \eta|_H,$$

flatly contradicting that C is sufficiently independent. $\square$

3.11. LEMMA. *There is a perfect sufficiently independent subset of $\hat{A}$.*

Proof. We will use Mycielski-Kuratowski's Theorem ([15], Theorem 19.1, p. 129.) For each infinite subgroup $H < A$ let

$$R_H = \{(\chi, \eta) \in \hat{A} \times \hat{A} : P_H(\chi) \neq P_H(\eta)\}$$

and for each sequence $i_1, \dots, i_n \in \mathbb{N}$ where such that $[A : H_{i_j}] < \infty$ for all $1 \leq j \leq n$ and $s : \{1, \dots, n\} \rightarrow \{-1, 1\}$ let

$$Q_s^{(i_1, \dots, i_n)} = \{(\chi_1, \dots, \chi_m, \eta) \in \hat{A}^{m+1} : \prod_{j=1}^n \chi_j^{s(j)} \circ \varphi_{i_j}|_{H_{(i_1, \dots, i_n)}} \neq \eta|_{H_{(i_1, \dots, i_n)}}\}$$

where

$$H_{(i_1, \dots, i_n)} = \bigcap_{j=1}^n H_{i_j}.$$

CLAIM 1. For each infinite $H < A$ the relation R_H is comeagre in $\hat{A} \times \hat{A}$.

Proof. By Kuratowski-Ulam's Theorem it suffices to show that for each $\eta \in \hat{A}$ the set

$$\{\chi \in \hat{A} : P_H(\chi) = P_H(\eta)\} = \eta\{\chi \in \hat{A} : \chi|H = 1\}$$

is meagre, and so to show that $\{\chi \in \hat{A} : \chi|H = 1\}$ is meagre. Suppose not. Then by Pettis' Theorem ([16], 9.9) it contains an open neighborhood of the identity. The map $\chi \mapsto \chi|H$ is open by the open mapping Theorem, and so it follows that $\{1\}$ is open in $\hat{H}$. Thus $\hat{H}$ must be discrete, and so H itself must be finite. $\square$ (Claim 1)

CLAIM 2. $Q_s^{(i_1, \dots, i_n)}$ is comeagre in $\hat{A}^{n+1}$.

Proof. Let $H = H_{(i_1, \dots, i_n)}$. By Kuratowski-Ulam's Theorem it is sufficient to show for each $\eta \in \hat{A}$ that the set

$$\{(\chi_1, \dots, \chi_n) \in \hat{A}^n : \prod_{j=1}^n \chi_j^{s(j)} \circ \varphi_{i_j}|H = \eta|H\}$$

is meagre. Define $\theta : \hat{A}^k \rightarrow \hat{H}$ by

$$\theta(\chi_1, \dots, \chi_k) = \prod_{j=1}^n \chi_j^{s(j)} \circ \varphi_{i_j}|H.$$

Then θ is a continuous homomorphism. In fact θ is onto since if $\chi \in \hat{H}$ then let $\tilde{\chi} \in \hat{A}$ be a character extending $\chi^{s(1)} \circ \varphi_{i_1}^{-1}$ to A , then

$$\theta(\tilde{\chi}, 1, \dots, 1) = \chi.$$

Thus by the open mapping Theorem θ is open, and so by Pettis' Theorem if the set

$$\ker(\theta) = \{(\chi_1, \dots, \chi_n) \in \hat{A}^n : \theta(\chi_1, \dots, \chi_n) = 1\}$$

is non-meagre then $\{1\}$ is open in $\hat{H}$, and so $\hat{H}$ must be discrete and H must be finite. But this contradicts that $[A : H] < \infty$. So $\ker(\theta)$ is meagre, and so since

$$(\eta_1, \dots, \eta_n) \ker(\theta) = \{(\chi_1, \dots, \chi_n) \in \hat{A}^n : \prod_{j=1}^n \chi_j^{s(j)} \circ \varphi_{i_j}|H = \eta|H\}$$

where $\theta(\eta_1, \dots, \eta_n) = \eta|H$, it follows that this set is meagre. $\square$ (Claim 2)

Applying the Mycielski-Kuratowski Theorem, it follows from Claim 1 and 2 that there is a perfect set $C \subseteq \hat{A}$ such that for all i such that H_i is infinite and j such that H'_j is infinite and all $\chi, \eta \in C$ distinct that $(\chi, \eta) \in R_{H_i}$ and $(\chi, \eta) \in R_{H'_j}$, and for all $\chi_1, \dots, \chi_{k+1} \in C$ distinct we have $(\chi_1, \dots, \chi_{k+1}) \in Q_s^{(i_1, \dots, i_n)}$. Then this C is a perfect sufficiently independent set. $\square$

Proof of Theorem 1. By Theorem 5.1 in [17], we have that $\approx^{P_c(2^{\mathbb{N}})}$ is a generically turbulent equivalence relation. Thus $E_0 <_{B \approx_c^{2^{\mathbb{N}}}}$ and $\approx_c$ is not classifiable by countable structures.

Let $C \subseteq \hat{A}$ be a perfect sufficiently independent subset, which we assume has Haar measure 0. For each $\sigma \in P_c(C)$ let $\hat{\sigma}$ be the symmetrization and let $\rho_{\hat{\sigma}}$ be defined as in 3.4. It is clear that if $\sigma, \sigma' \in P_c(C)$ are absolutely equivalent then $\rho_{\hat{\sigma}}$ and $\rho_{\hat{\sigma}'}$ are conjugate. If on the other hand $\sigma \not\approx \sigma'$, then by Lemma 3.10 $\rho_{\hat{\sigma}}|A$ and $\rho_{\hat{\sigma}'}|A$ are not conjugate, since otherwise $\Phi_{\hat{\sigma}} \approx \Phi_{\hat{\sigma}'}$.

If $\rho_{\hat{\sigma}}$ is not a.e. free, then we may instead consider the product with the Bernoulli shift action on $(\mathbb{R}^G, \mu_{\varphi_{reg}})$, where $\mu_{\varphi_{reg}}$ is the product of standard $N(0, 1)$ Gaussian probability measures. Then this action is free and as an A -action it has same spectral properties as $\rho_{\hat{\sigma}}$.

Since the assignment $P_c(C) \rightarrow P_c(\mathbb{R}^G) : \sigma \mapsto \mu_{\sigma}$ is continuous we have that

$$\approx^{P_c(C)} \leq_B (\simeq^{\mathcal{W}_A^*(G, X, \mu)}, \simeq_A^{\mathcal{W}_A^*(G, X, \mu)}).$$

Since for $A \leq H \leq G$ we have that $\simeq^{\mathcal{W}_A^*(G, X, \mu)} \subseteq \simeq_H^{\mathcal{W}_A^*(G, X, \mu)} \subseteq \simeq_A^{\mathcal{W}_A^*(G, X, \mu)}$ the remaining claims of Theorem 1 now follow from Proposition 2.4 and [17] Theorem 5.1. $\square$

3.12. The not type I case. The argument above unfortunately depends completely on the fact that Abelian groups have a “nice” representation theory. In case H is not abelian by finite, i.e. not type I, it is known by a result of Hjorth [12] that the infinite dimensional irreducible unitary representations of H cannot be classified by countable structures, in fact unitary equivalence $\sim_u$ in the space $\text{Irr}_{\infty}(H)$ of infinite dimensional irreducible representations has a turbulent part. Thus we have the following:

THEOREM 2. *Suppose G is a countable group and H is an infinite subgroup which is not abelian-by-finite. Let (X, μ) be a non-atomic standard Borel probability space, and let $\mathcal{X}$ be an infinite dimensional separable*

Hilbert space. Then $\sim_u^{\text{Irr}\infty(H, \mathcal{X})} \leq_B^\omega \simeq_H^{\mathcal{W}^*(G, X)}$. So in particular $\simeq_H^{\mathcal{W}^*(G, X)}$ is not classifiable by countable structures and $E_0 <_B \simeq_H^{\mathcal{W}^*(G, X)}$

Proof. (The proof follows the idea in [23], p. 110-111. See also Appendix E of [16] for a detailed account of this argument.) Let π be an irreducible unitary representation of H on $\mathcal{X}$. Then we can find a weakly mixing Gaussian action ρ_π of G such that $\text{Ind}_H^G(\pi)$ is a summand of the Koopman representation κ . By Theorem E.1 in [16], $\pi \mapsto \rho_\pi$ maps equivalent unitary representation π to conjugate G -actions. On the other hand, since $\pi < \kappa|_H$ at most countably many inequivalent π can give rise to conjugate-on- H actions. Thus $\pi \mapsto \rho_\pi$ is a countable-to-1 reduction of $\sim_u$ to $\simeq_H$, and the Theorem follows from Proposition 2.4. $\square$

Unfortunately the above actions may fail to be ergodic on $H \leq G$, since in general $\text{Ind}_H^G(\pi)|_H$ may not be weakly mixing. If, however, $H \triangleleft G$ is a normal infinite subgroup and $(\pi, \mathcal{X})$ is a weakly mixing representation of H , then $\text{Ind}_H^G(\pi)|_H$ is weakly mixing. More generally, we will say that a subgroup $H \leq G$ is *index stable* if for all $g \in G$ we have that

$$[H : gHg^{-1} \cap H] < \infty \iff [H : g^{-1}Hg \cap H] < \infty.$$

Then we have:

THEOREM 3. *Suppose $H < G$ is an infinite index stable subgroup of G which is not abelian by finite. If $\mathcal{X}$ is an infinite dimensional separable Hilbert space then*

$$\sim_u^{\text{Irr}\infty(H, \mathcal{X})} \leq_B^\omega \simeq_H^{\mathcal{W}_H^*(G, X)}$$

Proof. It suffices to show that if $(\pi, \mathcal{X})$ is a weakly mixing unitary representation of H then $\text{Ind}_H^G(\pi)|_H$ is weakly mixing.

We again identify $\text{Ind}_H^G(\pi)$ with π^α for some cocycle $\alpha : G \times G/H \rightarrow H$ as defined in [16], appendix G. Let $f : G/H \rightarrow \mathcal{X}$ be a unit vector in $l^2(G/H, \mathcal{X})$. We claim that $\langle \pi^\alpha(h)(f) : h \in H \rangle$ is not finite dimensional.

For this, suppose $f(gH) \neq 0$ for some $g = g_i$. Assume first that $[H : g^{-1}Hg \cap H] < \infty$. Since for $h \in gHg^{-1} \cap H$ we have $hg = gh'$ for some $h' \in H$ we have $\alpha(h, gH) = g^{-1}hg$ for all $h \in gHg^{-1} \cap H$. So for $h \in gHg^{-1} \cap H$ we get

$$\pi^\alpha(h)(f)(gH) = \pi(\alpha(h^{-1}, gH)^{-1})(f(h^{-1}gH)) = \pi(g^{-1}hg)(f(gH)).$$

Let $P_{gH} : l^2(G/H, \mathcal{X}) \rightarrow \mathcal{X}$ be the projection map

$$P_{gH}(f) = f(gH).$$

Then the above shows that

$$P_{gH}(\langle \pi^\alpha(h)(f) : h \in gHg^{-1} \cap H \rangle) = \langle \pi(h)(P_{gH}(f)) : h \in g^{-1}Hg \cap H \rangle.$$

But the latter cannot be finite dimensional since that would contradict the Lemma 4.4, since we assumed that $[H : g^{-1}Hg \cap H] < \infty$.

So we may assume that $[H : g^{-1}Hg \cap H] = \infty$ for all cosets gH where $f(gH) \neq 0$. Suppose for a contradiction that $\langle \pi^\alpha(H)(f) \rangle = \langle \pi^\alpha(h)(f) : h \in H \rangle$ is finite dimensional, and let $\xi_1, \dots, \xi_n$ be an orthonormal basis for it. If $u \in \langle \pi^\alpha(H)(f) \rangle$ is a unit vector and

$$u = a_1\xi_1 + \dots + a_n\xi_n$$

then

$$\begin{aligned} \|P_{gH}(u)\| &\leq |a_1|\|P_{gH}(\xi_1)\| + \dots + |a_n|\|P_{gH}(\xi_n)\| \\ &\leq (|a_1| + \dots + |a_n|) \max_{1 \leq k \leq n} \|P_{gH}(\xi_k)\| \\ &\leq n \max_{1 \leq k \leq n} \|P_{gH}(\xi_k)\|. \end{aligned}$$

Let $r = \max_i \|f(g_i H)\|$ and let g_{i_0} be such that $\|f(g_{i_0} H)\| = r$. We have $r > 0$ since f is non-zero. Let

$$\Delta = \{g_i : (\exists k \leq n) \|P_{g_i H}(\xi_k)\| \geq r/n\}.$$

Then Δ is finite. It follows from the above that for $g_i \notin \Delta$ and unit vectors $u \in \langle \pi^\alpha(H)(f) \rangle$ that

$$\|P_{g_i H}(u)\| < r.$$

Now we must have that $h \mapsto hg_{i_0}H$ ranges over infinitely many cosets: Indeed, if $hg_{i_0}H = h'g_{i_0}H$ then we have $hg_{i_0}\tilde{h} = h'g_{i_0}$ for some $\tilde{h} \in H$ and so $h^{-1}h' = g_{i_0}\tilde{h}g_{i_0}^{-1}$. Thus

$$h^{-1}h' \in g_{i_0}Hg_{i_0}^{-1} \cap H.$$

So if $h, h' \in H$ are not in the same coset of $H/(g_{i_0}Hg_{i_0}^{-1} \cap H)$ then $hg_{i_0}H \neq h'g_{i_0}H$, which proves that $h \mapsto hg_{i_0}H$ has infinite range since we assumed that $[H : g_{i_0}^{-1}Hg_{i_0} \cap H] = \infty$, and this implies that $[H : g_{i_0}Hg_{i_0}^{-1} \cap H] = \infty$.

It follows that we can find $h \in H$ such that $hg_{i_0}H \neq g_iH$ for all $g_i \in \Delta$. But now for this h we have

$$\|\pi^\alpha(h)(f)(hg_{i_0}H)\| = \|\pi(\alpha(h, hg_{i_0})^{-1})(f(g_{i_0}H))\| = \|f(g_{i_0}H)\| = r$$

while by the above we also have

$$\|\pi^\alpha(h)(f)(hg_{i_0}H)\| = \|P_{hg_{i_0}H}(\pi^\alpha(h)(f))\| < r,$$

since $\pi^\alpha(h)(f)$ is a unit vector in $\langle \pi^\alpha(H)(f) \rangle$ and $hg_{i_0} \notin \Delta$. $\square$

§4. ORBIT EQUIVALENCE.

We will now prove an immediate generalization of Corollary 2.6 in [13].

4.1. LEMMA. *Let G be a countable group with the relative property (T) over the infinite subgroup $H < G$. Then at most countably many ergodic-on- H probability measure preserving actions of G that are not conjugate on H can be orbit equivalent.*

4.2. Before the proof, recall that if E is a measure preserving countable equivalence relation on a standard Borel probability space (X, μ) then the *inner group* or *full group* of E is

$$[E] = \{T \in \text{Aut}(X, \mu) : (\forall^\mu x) xET(x)\}.$$

Since E is countable, $[E]$ becomes a Polish group when given the *uniform* topology, which is induced by the metric

$$d_U(T, S) = \mu\{x \in X : T(x) \neq S(x)\}.$$

Further, we may define a Borel measure M on E by

$$M(A) = \int |A_x| d\mu(x).$$

Where $A_x = \{y : (x, y) \in A\}$. Since E is measure preserving

$$M(A) = \int |A^y| d\mu(y)$$

defines the same measure, where $A^y = \{x : (x, y) \in A\}$. The equivalent integral form is

$$\int f(x, y) dM(x, y) = \int \sum_{y \in [x]_E} f(x, y) d\mu(x).$$

For each $T \in [E]$ there is a corresponding unit vector in $L^2(E)$,

$$\Delta_T(x, y) = \begin{cases} 1 & \text{if } T(x) = y \\ 0 & \text{if } T(x) \neq y \end{cases}$$

and so it is natural to think of the space $L^2(E, M)$ as a “linearization” of $[E]$.

Proof of Lemma 4.1. Let (Q, ε) be a Kazhdan pair for (G, H) such that any (Q, ε) -invariant unit vector is within $\frac{1}{10}$ of an H -invariant vector. Let $\sigma \in \mathcal{E}_H(G, X, \mu)$ and denote by $\mathcal{A}_\sigma$ the set of $\tau \in \mathcal{E}_H(G, X, \mu)$ such that $E_\tau \subseteq E_\sigma$ (a.e.). We equip $\mathcal{A}_\sigma$ with the topology it inherits as a subspace of $[E_\sigma]^G$ when $[E_\sigma]$ is given the uniform topology. Note that $\mathcal{A}_\sigma$ is separable.

We will show that conjugacy on H in $\mathcal{A}_\sigma$ is an open equivalence relation. Then it follows by separability that $\mathcal{A}_\sigma$ can only meet countably many $\sim_H$ classes, from which the Lemma follows.

Let $\tau_1, \tau_2 \in \mathcal{A}_\sigma$ and suppose for all $g \in Q$ that

$$\mu(\{x \in X : \tau_1(g)^{-1}(x) \neq \tau_2(g)^{-1}(x)\}) < \frac{\varepsilon^2}{2}.$$

We claim that τ_1 and τ_2 are conjugate on H . To see this, define a unitary representation π^{τ_1, τ_2} on $L^2(E_\sigma, M)$ by

$$\pi^{\tau_1, \tau_2}(g)(f)(x, y) = f(\tau_1(g)^{-1}(x), \tau_2(g)^{-1}(y)).$$

Denote by I the identity transformation. Then clearly $\pi^{\tau_1, \tau_2}(g)(\Delta_I) = \Delta_{\tau_2(g)\tau_1(g)^{-1}}$, and so for $g \in Q$ we have from our assumption that

$$\begin{aligned} \|\pi^{\tau_1, \tau_2}(g)(\Delta_I) - \Delta_I\|_{L^2(E_\sigma)}^2 &= \|\Delta_{\tau_2(g)\tau_1(g)^{-1}} - \Delta_I\|_{L^2(E_\sigma)}^2 \\ &= \int \sum_{y \in [x]_{E_\sigma}} |\Delta_{\tau_2(g)\tau_1(g)^{-1}}(x, y) - \Delta_I(x, y)|^2 d\mu(x, y) < \varepsilon^2. \end{aligned}$$

Thus Δ_I is (Q, ε) -invariant, and so there is a H -invariant vector $f \in L^2(E_\sigma)$ with $\|f - \Delta_I\|_{L^2(E_\sigma)} \leq \frac{1}{10}$. Since $\|f - \Delta_I\|_{L^2(E_\sigma)} \leq \frac{1}{10}$ we must have that

$$\mu(\{x \in X : |1 - f(x, x)| \geq \frac{1}{2}\}) < \frac{1}{4}.$$

On the other hand, we must also have that

$$\mu(\{x \in X : (\exists y_1, y_2) y_1 \neq y_2 \wedge f(x, y_1) > \frac{1}{2} \wedge f(x, y_2) > \frac{1}{2}\}) < \frac{1}{4}.$$

It follows that the set

$$A = \{x \in X : (\exists! y) f(x, y) > \frac{1}{2}\}$$

has $\mu(A) > 0$. Since moreover f is H -invariant, A is $\tau_1|H$ -invariant. Since τ_1 is ergodic in H we must have $\mu(A) = 1$.

Define $\varphi(x)$ to be the unique $y \in X$ such that $f(x, y) > \frac{1}{2}$. Then clearly $\varphi \subseteq E_\sigma$. It is also clear from the above argument that φ is invertible, since τ_2 is ergodic on H . Thus φ is measure preserving.

From the H -invariance of f we now get that for $h \in H$,

$$\frac{1}{2} < f(x, \varphi(x)) = \pi^{\tau_1, \tau_2}(h)(f)(x, \varphi(x)) = f(\tau_1(h)^{-1}(x), \tau_2(h)^{-1}(\varphi(x))).$$

Since $\varphi(x)$ is the unique y where $f(x, y) > 1/2$ it follows that

$$\varphi(\tau_1(h)^{-1}(x)) = \tau_2(h)^{-1}(\varphi(x))$$

for all $h \in H$. Thus φ witnesses that τ_1 and τ_2 are conjugate on H . $\square$

Let (X, μ) be a standard Borel probability space. We denote by $\simeq_{OE}$ the equivalence relation of orbit equivalence in $\mathcal{A}(G, X, \mu)$. Combining Lemma 4.1 and Theorem 1 and 3, we then have:

THEOREM 4. *Let G be a countable group with the relative property (T) over an infinite subgroup H , and let (X, μ) be a standard Borel probability space. If either*

(i) *H is Abelian or*

(ii) *H is not Abelian-by-finite and is index stable in G ,*

then G has continuum many orbit inequivalent weakly mixing a.e. free actions on (X, μ) . In fact, $E_0 <_B \simeq_{OE}^{\mathcal{W}^(G, X, \mu)}$ and $\simeq_{OE}^{\mathcal{W}^*(G, X, \mu)}$ is not classifiable by countable structures.*

Proof. (i) By Theorem 1 we have $\approx^{P_c(2^{\mathbb{N}})} \leq_B (\simeq_{\mathcal{W}_A^*(G, X, \mu)}, \simeq_A^{\mathcal{W}_A^*(G, X, \mu)})$. Let $\theta : P_c(2^{\mathbb{N}}) \rightarrow \mathcal{W}_A^*(G, X, \mu)$ witness this. Then θ is also homomorphism of $P_c(2^{\mathbb{N}})$ into $\simeq_{OE}^{\mathcal{W}^*(G, X, \mu)}$. Since if $\mu, \mu' \in P_c(2^{\mathbb{N}})$ are absolutely inequivalent measures then $\theta(\mu)$ and $\theta(\mu')$ are not conjugate on A , it follows by Lemma 4.1 that at most countably many absolutely inequivalent measures in $P_c(2^{\mathbb{N}})$

can give rise to orbit equivalent actions. Thus θ is Borel countable-to-1 reduction of absolute equivalence in $P_c(2^{\mathbb{N}})$ to $\simeq_{OE}^{\mathcal{W}^*(G,X,\mu)}$. Theorem 4 (i) now follows from Proposition 2.4 and Theorem 5.1 in [17].

(ii) Let $\mathcal{H}$ be an infinite dimensional Hilbert space. It is clear that the reduction $\pi \mapsto \rho_\pi$ given in Theorem 2 is a homomorphism of $\sim_u$ in $\text{Irr}_\infty(H, \mathcal{H})$ to $\simeq$ and therefore also to $\simeq_{OE}$. If H is index stable then by Theorem 3 the action ρ_π are weakly mixing on H . By Lemma 4.1 it follows that only countably many inequivalent π can give rise to ρ_π that are conjugate on H . Therefore the homomorphism $\pi \mapsto \rho_\pi$ is countable-to-1 reduction of $\sim_u^{\text{Irr}_\infty(H)}$ to $\simeq_{OE}^{\mathcal{W}^*(G,X,\mu)}$. $\square$

References

- [1] H. Becker, A. Kechris, *The descriptive set theory of Polish group actions*, London Mathematical Society lecture notes, vol. 232 (1996), Cambridge University Press.
- [2] B. Bekka, P. de la Harpe, A. Valette, *Kazhdan's Property (T)*, to appear, (2003).
- [3] V.I. Bogachev, *Gaussian Measures*, Mathematical Surveys and Monographs, American Mathematical Society (1998).
- [4] A. Connes, J. Feldman, B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergodic Theory and Dynamical Systems, vol. 1(1981), pp. 431–450.
- [5] A. Connes, B. Weiss, *Property T and asymptotically invariant sequences*, Israel Journal of Mathematics, vol. 37 (1980), pp. 209–210.
- [6] H.A. Dye, *On groups of measure preserving transformation. I*, American Journal of Mathematics, vol. 81(1959) pp. 119–159.
- [7] H.A. Dye, *On groups of measure preserving transformation. II*, American Journal of Mathematics, vol. 85(1963) pp. 551–576.
- [8] M. Foreman, B. Weiss, *An anti-classification theorem for ergodic measure preserving transformations*. J. Eur. Math. Soc. 6 (2004), no. 3, pp. 277–292
- [9] Glasner, E., *Ergodic Theory via Joinings*, Mathematical Surveys and Monographs no. 101, American Mathematical Society (2003).
- [10] E. Glasner, B. Weiss *Kazhdan's property T and the geometry of the collection of invariant measures*, GAFA, Geom. Funct. anal. 7, (1997), pp. 917–935.
- [11] P. Halmos, *Lectures on ergodic theory*, Chelsea Publishing Co., New York (1956).

- [12] G. Hjorth, *When it's bad it's worse*, (note), (1997).
- [13] G. Hjorth, *A converse to Dye's theorem*, Trans. Amer. Math. Soc., 357 (2005) no. 8, pp. 3083–3103
- [14] G. Hjorth, *Classification and Orbit Equivalence Relations*, Mathematical Surveys and Monographs 75, American Mathematical Society (2000)
- [15] A. Kechris, *Classical descriptive set theory*, Springer Verlag, New York, (1995).
- [16] A. Kechris, *Global aspects of ergodic group actions and equivalence relations*, preprint, Caltech, (2006).
- [17] A. Kechris, N.E. Sofronidis, *A strong ergodicity property of unitary and self-adjoint operators*, Ergodic Theory and Dynamical Systems 21, pp. 1459–1479, (2001).
- [18] D. Ornstein, B. Weiss, *Ergodic theory of amenable group actions*, Bulletin of the American Mathematical Society, vol. 2(1980), pp. 161–164.
- [19] Popa, S. *Some computations of 1-cohomology groups and construction of non-orbit-equivalent actions*, J. Inst. Math. Jussieu 5 (2006), no. 2, pp. 309–332
- [20] I.P. Cornfield, S.V. Fomin, Ya. G. Sinai, *Ergodic Theory*, Springer-Verlag, New York, (1982).
- [21] K. Schmidt, *Asymptotically invariant sequences and an action of $SL(2, \mathbb{Z})$ on the sphere*, Israel Journal of Mathematics, vol. 37 (1980), pp. 193–208.
- [22] A. Törnquist, *The Borel complexity of orbit equivalence*, Ph.D. Thesis, UCLA (2005).
- [23] R. Zimmer, *Ergodic theory and semisimple Lie groups*, Birkhäuser, (1984).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO
 40 ST. GEORGE STREET, ROOM 6092, TORONTO, ONTARIO, CANADA
E-mail: asger@math.utoronto.ca